

AN OPERADIC MODEL FOR A MAPPING SPACE AND ITS ASSOCIATED SPECTRAL SEQUENCE

DAVID CHATAUR AND KATSUHIKO KURIBAYASHI

ABSTRACT. Let X and Y be simplicial sets and $\overline{\mathbb{F}}_p$ the closure of the finite field \mathbb{F}_p . In [11], we give an algebra model over an $E_\infty\overline{\mathbb{F}}_p$ -operad \mathcal{E} for the mapping space $\mathcal{F}(X, Y)$, whose source X is finite, provided the connectivity of Y is greater than or equal to $\dim X$. Moreover, the model enables us to construct a spectral sequence converging to the cohomology of $\mathcal{F}(X, Y)$ with coefficients in $\overline{\mathbb{F}}_p$, whose E_2 -term is expressed via Lannes' division functor in the category of unstable $\overline{\mathbb{F}}_p$ -algebra over the Steenrod algebra. In this article, we survey important results described in [11].

1. OVERVIEW

Let X and Y be spaces (or simplicial sets) and $\mathcal{F}(X, Y)$ denote the mapping space. In [13], Haefliger has given a rational model for a mapping space $\mathcal{F}(X, Y)$ for which Y is a nilpotent space. Subsequently, Bousfield, Peterson and Smith [5] have constructed another rational model for a mapping space with a functorial way, more precisely, their model is obtained by using a division functor in the category of commutative differential \mathbb{Z} graded algebras over the rational field. In the same paper, we are also aware of an interesting spectral sequence (henceforth BPS spectral sequence) converging to $H^*(\mathcal{F}(X, Y); \mathbb{Q})$, which is constructed with the algebraic model. Brown and Szczarba [6] have derived an accessible rational model for $\mathcal{F}(X, Y)$ by computing the division functor explicitly. The construction renders the model more computable. For more recent progress of the material, we refer the reader to [7], [8], [14], [18] and [19].

As for a p -adic model for a space, Mandell [20] has proved that the homotopy category of nilpotent, p -complete spaces of finite p -type is equivalent to a full subcategory of the homotopy category of algebras over an $\overline{\mathbb{F}}_p$ -operad \mathcal{E} . Here $\overline{\mathbb{F}}_p$ denotes the closure of the finite field \mathbb{F}_p . This motivates us to construct an \mathcal{E} -algebra model for a mapping space $\mathcal{F}(X, Y)$. Recently, Fresse [12] has given such a model by means of a division functor in the category of algebras over an $\overline{\mathbb{F}}_p$ -operad under some finiteness condition on the homotopy group of X . One of the purposes of the paper [11] is to improve Fresse's model for a mapping space. Another one is to construct a spectral sequence converging to $H^*(\mathcal{F}(X, Y); \overline{\mathbb{F}}_p)$, which is regarded as a p -adic version of the BPS spectral sequence.

It is worth to mention that Chataur and Thomas [10] have constructed a \mathcal{C}_∞ -algebra model for a mapping space which is the normalization of a simplicial

2000 Mathematics Subject Classification. 55P48, 18D50.

Key words and phrases. algebras over an operad, the division functor, mapping spaces.

Département de Mathématiques, Université des Sciences et Technologies de Lille, Cité Scientifique - Bâiment M2, F-59655 Villeneuve d'Ascq Cedex, France David.Chataur@math.univ-lille1.fr

Department of Mathematical Sciences, Faculty of Science, Shinshu University, Matsumoto, Nagano 390-8621, Japan kuri@math.shinshu-u.ac.jp

\mathcal{C}_∞ -algebra, where \mathcal{C}_∞ is a cofibrant replacement of the commutative operad. When the source space is the circle, the model for the mapping space, namely for the loop space, is related with the Hochschild chain complex. We also mention that our spectral sequence is a generalization of that due to Bökstedt and Ottosen [4], which converges to the cohomology of a free loop space.

We recall briefly the algebraic model for a mapping space over an operad due to Fresse. Let \mathcal{E} denote the Barratt-Eccles operad over a field \mathbb{K} , which is an E_∞ -operad. Then we can regard the normalized cochain functor $C^*(-; \mathbb{K})$ as a functor from the category of simplicial sets to $\mathcal{E}\text{-Alg}$ the category of \mathcal{E} -algebras ([2, 1.5], [21]). Let A be an \mathcal{E} -algebra and K an \mathcal{E} -coalgebra. The diagonal map on \mathcal{E} makes the dg-module $\text{Hom}_{\mathbb{K}}(K, -)$ of homogeneous morphisms into an \mathcal{E} -algebra (see [12, 1.5] for details). We denote by $\mathcal{E}(,)$ the hom set in $\mathcal{E}\text{-Alg}$.

Proposition 1.1. [12, 1.6.Proposition] *Let K be an \mathcal{E} -coalgebra. Then the functor $\text{Hom}_{\mathbb{K}}(K, -)$ has a left adjoint. More explicitly, for A an \mathcal{E} -algebra, there is an \mathcal{E} -algebra $A \otimes K$ such that $\mathcal{E}(A \otimes K, -) \cong \mathcal{E}(A, \text{Hom}_{\mathbb{K}}(K, -))$.*

Let K^* be an \mathcal{E} -algebra of finite type and K_* the \mathcal{E} -coalgebra which is the dual to K^* . Then, by definition, $A \otimes K_*$ is regarded as Lannes' functor $(A : K^*)_{\mathcal{E}\text{-Alg}}$ in the category of \mathcal{E} -algebras (see [22, 3.2 and 3.8] for the existence of the division functor, such as Lannes' T -functor). Moreover, if A is an almost free algebra $\mathcal{E}(V)$, then $A \otimes K$ is also an almost free algebra of the form $\mathcal{E}(V \otimes K)$. Since $\text{Hom}_{\mathbb{K}}(K, -)$ preserves fibrations and acyclic fibrations, the total left derived functor $- \otimes^L K$ of $- \otimes K$ can be defined; that is, we have a natural bijection $\bar{h}\mathcal{E}(A \otimes^L K, -) \cong \bar{h}\mathcal{E}(A, \text{Hom}_{\mathbb{K}}(K, -))$ for any \mathcal{E} -algebra A . Here $\bar{h}\mathcal{E}(,)$ denotes the hom set in the homotopy category of \mathcal{E} -algebras. The functor $- \otimes^L K$ provides an \mathcal{E} -algebra model for a mapping space.

Theorem 1.2. [12, 1.10.Theorem] *Let X and Y be simplicial sets. We assume that X is finite and that $\pi_n(Y)$ is a finite p -group for $n \geq 0$. We have a quasi-isomorphism between $C^*(\mathcal{F}(X, Y); \mathbb{K})$ and $C^*(Y; \mathbb{K}) \otimes^L C_*(X; \mathbb{K})$, which is functorial with respect to X and Y .*

Henceforth, we work in the category of algebras over the Barratt-Eccles operad \mathcal{E} defined in the field $\overline{\mathbb{F}}_p$. The chain and cochain complexes $C_*(X; \overline{\mathbb{F}}_p)$ and $C^*(X; \overline{\mathbb{F}}_p)$ are written as $C_*(X)$ and $C^*(X)$, respectively. In [11], we first show that $C^*(\mathcal{F}(X, Y))$ can be connected with $C^*(Y) \otimes^L C_*(X)$ by quasi-isomorphisms without assuming that $\pi_n(Y)$ is a finite p -group, subject to the connectedness of the mapping space $\mathcal{F}(X, Y)$. More precisely, the following theorem is established.

Theorem 1.3. [11, Theorem 1.3] *Let X be a finite simplicial set and Y a connected nilpotent simplicial set of finite type. Assume that the connectivity of Y is greater than or equal to the dimension of X . Then there exists an isomorphism between $C^*(\mathcal{F}(X, Y))$ and $C^*(Y) \otimes^L C_*(X)$, which is functorial with respect to X and Y , in the homotopy category of \mathcal{E} -algebras.*

As mentioned above, the functor $- \otimes K_*$ is regarded as Lannes' division functor $(- : K^*)_{\mathcal{E}\text{-Alg}}$. This fact enables us to construct a spectral sequence converging to the cohomology $H^*(\mathcal{F}(X, Y))$. In order to describe the spectral sequence more precisely, we recall that the generalized Steenrod algebra \mathcal{B} is the free associative $\overline{\mathbb{F}}_p$ -algebra generated by the P^s and (if $p > 2$) the βB^s for $s \in \mathbb{Z}$ over the two

sided ideal generated by the Adem relations (see [20, Section 11]). The result [20, Theorem 1.4] states that the quotient algebra $\mathcal{B}/(Id - P^0)$ is the usual Steenrod algebra \mathcal{A} . Let $\mathcal{K}\text{-}\overline{\mathbb{F}}_p$ be the category of unstable $\overline{\mathbb{F}}_p$ -algebras over the generalized Steenrod algebra \mathcal{B} . We have a spectral sequence.

Theorem 1.4. [11, Theorem 1.4] (Compare with [5, Corollary 3.5]) *Let X be a finite simplicial set and Y a connected nilpotent simplicial set of finite type. Assume that the connectivity of Y is greater than or equal to the dimension of X . Then there exists a left-half plane spectral sequence $\{E_r, d_r\}$ with*

$$E_2^{s,*} \cong L_s(H^*(Y) : H^*(X))_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p}$$

converging strongly to $H^(\mathcal{F}(X, Y))$. Here $L_s(- : H^*(X))_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p}$ denotes the s^{th} left derived functor of the division functor $(- : H^*(X))_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p}$ in the category $\mathcal{K}\text{-}\overline{\mathbb{F}}_p$. Moreover the spectral sequence is natural with respect to X and Y .*

In what follows, we shall refer to the spectral sequence in Theorem 1.4 as the mod p BPS spectral sequence. For a \mathcal{B} -algebra B and a \mathcal{B} -algebra A of finite type, one can define the derived functor $L_s(B : A)_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p}$ using a simplicial resolution of B in the category $\mathcal{K}\text{-}\overline{\mathbb{F}}_p$. Since the resolution is a complex in the category of unstable \mathcal{B} -modules, the functor $L_s(B : A)_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p}$ for any s inherits the \mathcal{B} -module structure from that of the complex. The same derived functor can be defined in the category $\mathcal{S}\text{-}\overline{\mathbb{F}}_p$ of unstable $\overline{\mathbb{F}}_p$ -algebras over the usual Steenrod algebra \mathcal{A} . Observe that an object in $\mathcal{S}\text{-}\overline{\mathbb{F}}_p$ is regarded as one in $\mathcal{K}\text{-}\overline{\mathbb{F}}_p$ with the natural projection $\mathcal{B} \rightarrow \mathcal{B}/(Id - P^0) = \mathcal{A}$. The following theorem allows us to work in the more familiar category $\mathcal{S}\text{-}\overline{\mathbb{F}}_p$ than $\mathcal{K}\text{-}\overline{\mathbb{F}}_p$ when computing the mod p BPS spectral sequence.

Theorem 1.5. [11, Theorem 1.5] *Let A and B be \mathcal{A} -algebras of finite type. Then $L_s(B : A)_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p}$ is isomorphic to $L_s(B : A)_{\mathcal{S}\text{-}\overline{\mathbb{F}}_p}$ as a \mathcal{B} -module for any s .*

This theorem implies that the mod p BPS spectral sequence is reducible in the second quadrant. Moreover we have

Assertion 1.6. *The mod p BPS spectral sequence possesses an unstable module structure on \mathcal{B} and hence on \mathcal{A} .*

For the more precise statement concerning the Steenrod operations on the spectral sequence, see [11, Theorem 7.7].

Unfortunately, we are less successful in computing the BPS spectral sequence. However we firmly believe the spectral sequence to be of use in the study of mapping spaces. As for the edge homomorphism, we have an interesting example (Theorem 1.7 below). Let Y be a simply connected space whose cohomology is a polynomial algebra. Then the results [15, Remarks 3.4, 3.5] and [17, Theorem 1.6] enable one to determined explicitly the mod p cohomology of the free loop space $LY = \mathcal{F}(S^1, Y)$. To be exact, if $H^*(Y; \mathbb{F}_p) = \mathbb{F}_p[y_1, \dots, y_l]$, as an $H^*(Y; \mathbb{F}_p)$ -algebra, then $H^*(LY; \mathbb{F}_p) \cong \mathbb{F}_p[y_1, \dots, y_l] \otimes \Lambda(\bar{y}_1, \bar{y}_2, \dots, \bar{y}_l)$ if $p \neq 2$, where $\deg \bar{y}_i = \deg y_i - 1$. In the case $p = 2$, we see that

$$H^*(LY; \mathbb{F}_2) \cong \mathbb{F}_2[y_1, \dots, y_l] \otimes \mathbb{F}_2[\bar{y}_1, \bar{y}_2, \dots, \bar{y}_l] / (\bar{y}_i^2 + \mathfrak{D}S q^{\deg y_i - 1} y_i; i = 1, 2, \dots, l)$$

as an $H^*(Y; \mathbb{F}_2)$ -algebra, for which \mathfrak{D} is the derivation defined by $\mathfrak{D}(y_i) = \bar{y}_i$

Since the derivation \mathfrak{D} is compatible with the Steenrod operations ([15, Remark 3.5]), we can determine explicitly the \mathcal{A} -algebra structure of $H^*(LY; \mathbb{F}_p)$ from that of the polynomial algebra $H^*(Y; \mathbb{F}_p)$ (see, for example, [15, Example 3.6]). The following theorem asserts that the \mathcal{A} -algebra structure of $H^*(LY; \overline{\mathbb{F}}_p)$ is also expressed via the Lannes' division functor.

Theorem 1.7. [11, Theorem 6.4] *Let Y be a simply-connected space whose mod p cohomology is a polynomial algebra. Then the edge homomorphism*

$$\text{edge}_{(Y; S^1)} : (H^*(Y) : H^*(S^1))_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p} \rightarrow H^*(LY)$$

is an isomorphism.

We conclude this note by describing briefly other topics in [11]. In Section 5, we prove that, for any unstable \mathcal{A} -algebra A of finite type, $L_s(H^*(K(\mathbb{Z}/p, n) : A))_{\mathcal{K}\text{-}\overline{\mathbb{F}}_p} = 0$ when $s < 0$. This fact is a key to proving Theorem 1.5. Section 7, Appendix, is devoted to defining well-behaved Steenrod operations in the spectral sequence which is constructed out of a simplicial \mathcal{E} -algebra. In consequence, not only does we can establish an *algebraic construction* of the Steenrod operations on the Eilenberg-Moore spectral sequence also we see that the mod p BPS spectral sequence possesses an unstable module structure on \mathcal{A} .

REFERENCES

- [1] M. Barr and J. Beck, Homology and standard constructions, in Seminar on triples and categorical homology theory, Springer Lecture Notes in Math. 80(1969), 245-335.
- [2] C. Berger and B. Fresse, Combinatorial operad actions on cochains, preprint [arXiv:math.AT/0109158](https://arxiv.org/abs/math/0109158) (2002).
- [3] M. Bökstedt and I. Ottosen, Homotopy orbits of free loop spaces, Fund. Math. 162 (1999), 251-275.
- [4] M. Bökstedt and I. Ottosen, A spectral sequence for string cohomology, preprint (2002).
- [5] A. K. Bousfield, C. Peterson and L. Smith, The rational homology of function spaces, Arch. Math. 52 (1989), 275-283
- [6] E. H. Brown Jr and R. H. Szczarba, Rational homotopy type of function spaces, Trans. Amer. Math. Soc. **349**(1997), 4931-4951.
- [7] U. Buijs and A. Murillo, Basic constructions in rational homotopy theory of function spaces, preprint (2005).
- [8] U. Buijs and A. Murillo, The rational homotopy Lie algebra of function spaces, preprint (2005).
- [9] D. Chataur, Formes différentielles généralisées sur une opérade et modèles algébriques des fibrations, Algebr. Geom. Topol. 2 (2002), 51–93.
- [10] D. Chataur and J.-C. Thomas, E_∞ -model for a mapping space, to appear in Topology Appl.
- [11] D. Chataur and K. Kuribayashi, An operadic model for a mapping space and its associated spectral sequence, preprint <http://marine.shinshu-u.ac.jp/kuri/dvi/E-algebra-neo3.pdf> (2005).
- [12] B. Fresse, Derived division functors and mapping spaces, preprint [arXiv:math.At/0208091](https://arxiv.org/abs/math/0208091) (2002). 1984–1986),
- [13] A. Haefliger, Rational homotopy of space of sections of a nilpotent bundle, Trans. Amer. Math. Soc. **207**(1982), 609-620.
- [14] Y. Hirato, K. Kuribayashi and N. Oda, A function space model approach to the rational evaluation subgroups, preprint (2005).
- [15] A. Kono and K. Kuribayashi, Module derivations and cohomological splitting of adjoint bundles, Fund. Math. 180 (2003), 199-221.
- [16] I. Kriz and J. P. May, Operads, algebras, modules and motives. Astérisque No. 233 (1995).
- [17] K. Kuribayashi, Module derivations and the adjoint action of a finite loop space, J. Math. Kyoto Univ. **39** (1999) 67-85.
- [18] K. Kuribayashi, A rational model for the evaluation map, preprint (2005).
- [19] K. Kuribayashi and T. Yamaguchi, A rational splitting of a based mapping space, preprint (2005).
- [20] M. Mandell, E_∞ algebras and p -adic homotopy theory. Topology 40 (2001), no. 1, 43–94.
- [21] J. E. McClure and J. H. Smith, Multivariable cochain operations and little n -cubes, J. Amer. Math. Soc. 16 (2003), 681-704.
- [22] L. Schwartz, Unstable modules over the Steenrod algebra and Sullivan's fixed point set conjecture, Chicago Lectures in Mathematics, University of Chicago Press, 1994.