

A double filtration for the mapping class group and the Goeritz group of the sphere

Anderson Vera (Joint work with K. Habiro)


Kyoto University/JSPS

Online Topology Seminar, The University of Tokyo

July 28, 2020


Mapping class group

$\Sigma = \Sigma_{g,1}$: Compact, connected, oriented surface of genus g with 1 boundary component.



$$\mathcal{M} = \{h : \Sigma \xrightarrow{\sim} \Sigma \mid h_{\partial\Sigma} = \text{Id}_{\partial\Sigma}\} / \text{isotopy}$$

Dehn twist:



Dehn-Nielsen representation

$$\pi = \pi_1(\Sigma, *) = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle$$

The map

$$\rho : \mathcal{M} \longrightarrow \text{Aut}(\pi), \quad h \longmapsto h_\#$$

is injective.

Lower central series

$$\Gamma_1\pi = \pi, \Gamma_2\pi = [\pi, \pi], \Gamma_3\pi = [\pi, [\pi, \pi]],$$

$$\Gamma_{n+1}\pi = [\pi, \Gamma_n\pi]$$

$$[a, b] = aba^{-1}b^{-1}$$

The action of \mathcal{M} on π preserves the lower central series.

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{\rho} & \text{Aut}(\pi) \\ & \searrow \rho_n & \downarrow \\ & & \text{Aut}(\pi/\Gamma_{n+1}\pi) \end{array}$$

Torelli group

$$\begin{aligned} \mathcal{I} &= \{h \in \mathcal{M} \mid h_* = \text{Id}_{H_1(\Sigma)}\} \\ &= \ker(\rho_1) \end{aligned}$$

Johnson filtration

$$\mathcal{I} = J_1 \mathcal{M} \supset J_2 \mathcal{M} \supset J_3 \mathcal{M} \supset \dots$$

$$\begin{aligned} J_n \mathcal{M} &= \ker(\rho_n) \\ &= \{h \in \mathcal{M} \mid \forall x \in \pi : h_{\#}(x)x^{-1} \in \Gamma_{n+1}\pi\} \\ &= \{h \in \mathcal{M} \mid [h, \pi] \subset \Gamma_{n+1}\pi\} \\ &\quad (\text{considering } \pi \rtimes \mathcal{M}) \end{aligned}$$

Underlying abstract setting: A group G (above $G = \mathcal{M}$) acting on a group K (above $K = \pi$) with a \mathbb{N}_+ -filtration $(K_i)_{i \geq 1}$ (above $K_i = \Gamma_i \pi$).


- (Habiro, Massuyeau) Considered a group G acting on a group K with a \mathbb{N} -filtration $(K_i)_{i \geq 0}$.
- We consider a group G acting on a group K with a Λ -filtration $(K_i)_{\lambda \in \Lambda}$. Where Λ is a good ordered commutative monoid

- In this talk we consider $\Lambda = \mathbb{N}^2 = \{(m, n) \in \mathbb{Z} \mid m, n \geq 0\}$

with order $(m, n) \leq (m', n')$ iff $m \leq m'$ and $n \leq n'$.

Goal


Define a **double filtration** for the mapping class group \mathcal{M} and for the Goeritz group \mathcal{G}



\mathbb{N}^2 -filtrations

A \mathbb{N}^2 -filtration of a group K is a family $(K_{m,n})_{(m,n) \in \mathbb{N}^2}$ of normal subgroups of K such that

- $K_{0,0} = K$
- $K_{m,n} \supset K_{m',n'} \text{ for } (m,n) \leq (m',n')$
- $[K_{m,n}, K_{m',n'}] \subset K_{m+m', n+n'} \text{ for all } (m,n), (m',n') \in \mathbb{N}^2$



$$\bar{K}_{m,n} := \frac{K_{m,n}}{K_{m+1,n} K_{m,n+1}}$$

is an **abelian group**.

Double lower central series

Consider a group K and two normal subgroups $\bar{X}, \bar{Y} \trianglelefteq K$. The double lower central series of (K, \bar{X}, \bar{Y}) is the \mathbb{N}^2 -filtration $(K_{m,n})_{(m,n) \in \mathbb{N}^2}$ of K given by

- $K_{0,0} = K$
- $K_{m,0} = \Gamma_m \bar{X}$ for $m \geq 1$
- $K_{0,n} = \Gamma_n \bar{Y}$ for $n \geq 1$
- $K_{m,n} = [K_{1,0}, K_{m-1,n}] [K_{0,1}, K_{m,n-1}]$
for $m, n \geq 1$

K	\supset	\bar{Y}	\supset	$\Gamma_2 \bar{Y}$	\supset	\dots
\cup		\cup		\cup		
\bar{X}	\supset	$K_{1,1}$	\supset	$K_{1,2}$	\supset	\dots
\cup		\cup		\cup		
$\Gamma_2 \bar{X}$	\supset	$K_{2,1}$	\supset	$K_{2,2}$	\supset	\dots
\cup		\cup		\cup		
$\Gamma_3 \bar{X}$	\supset	$K_{3,1}$	\supset	$K_{3,2}$	\supset	\dots
\cup		\cup		\cup		
\vdots		\vdots		\vdots		

$$K_{1,1} = [\bar{X}, \bar{Y}]$$

$$K_{2,1} = [\bar{X}, [\bar{X}, \bar{Y}]]$$

$$K_{2,2} = [\bar{Y}, [\bar{X}, [\bar{X}, \bar{Y}]]] [[\bar{X}, \bar{Y}], [\bar{X}, \bar{Y}]]$$

Lemma

Let K a group and $\bar{X}, \bar{Y} \trianglelefteq K$ such that $K = \bar{X}\bar{Y}$. Let $(K_{i,j})_{(i,j) \in \mathbb{N}^2}$ be the double lower central series of $(K; \bar{X}, \bar{Y})$. For $m \geq 1$, we have

$$\Gamma_m(K) = \prod_{i+j=m} K_{i,j}.$$

- $\Gamma_1 K = \bar{X}\bar{Y}$
- $\Gamma_2 K = K_{2,0} K_{1,1} K_{0,2}$
- $\Gamma_3 K = K_{3,0} K_{2,1} K_{1,2} K_{0,3}$

$$\begin{array}{ccccccc} K = \bar{X}\bar{Y} & \supset & \bar{Y} & \supset & \Gamma_2 \bar{Y} & \supset & \dots \\ \cup & & \cup & & \cup & & \\ \bar{X} & \supset & K_{1,1} & \supset & K_{1,2} & \supset & \dots \\ \cup & & \cup & & \cup & & \\ \Gamma_2 \bar{X} & \supset & K_{2,1} & \supset & K_{2,2} & \supset & \dots \\ \cup & & \cup & & \cup & & \\ \Gamma_3 \bar{X} & \supset & K_{3,1} & \supset & K_{3,2} & \supset & \dots \\ \vdots & & \vdots & & \vdots & & \end{array}$$

Goeritz group of \mathbb{S}^3

In term of basis:

- $K = \pi_1(\Sigma, *) = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle$ free group on $2g$ generators
- $\bar{X} = \langle\langle x_1, \dots, x_g \rangle\rangle_K$ normal closure in K
- $\bar{Y} = \langle\langle y_1, \dots, y_g \rangle\rangle_K$

Let $(K_{i,j})_{(i,j) \in \mathbb{N}^2}$ be the **double lower central series** of $(K; \bar{X}, \bar{Y})$.

The subgroup of the mapping class group

$$\mathcal{G} = \mathcal{G}_{g,1} = \{ h \in \mathcal{M} \mid h_{\#}(\bar{X}) \subset \bar{X}, \quad h_{\#}(\bar{Y}) \subset \bar{Y} \}$$

acts on K preserving the double lower central series of $(K; \bar{X}, \bar{Y})$

The group \mathcal{G} is called the (genus g) **Goeritz group of \mathbb{S}^3** (relative to the disk D)

Goeritz group of \mathbb{S}^3

$$K = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle \quad \bar{X} = \langle\langle x_1, \dots, x_g \rangle\rangle_K \quad \bar{Y} = \langle\langle y_1, \dots, y_g \rangle\rangle_K$$

$$\mathcal{G} = \{ h \in \mathcal{M} \mid h_{\#}(\bar{X}) \subset \bar{X}, \quad h_{\#}(\bar{Y}) \subset \bar{Y} \}$$

Equivalent definitions:

- \mathcal{G} is the group of isotopy classes of orientation-preserving homeomorphism $h : \mathbb{S}^3 \rightarrow \mathbb{S}^3$ such that $h(\partial V) = \partial V$ and $h(D) = D$.
- \mathcal{G} is the subgroup of mapping classes in \mathcal{M} which extend to both handlebodies V and V'
- \mathcal{G} is the subgroup of mapping classes in \mathcal{M} which preserve the standard Heegaard splitting of the 3-sphere.

Remark. It is not known if \mathcal{G} is finitely generated for genus > 3 .

Powell Conjecture. The group \mathcal{G} is finitely generated. Moreover, Powell proposed a set of generators with 5 elements.

Goeritz group of \mathbb{S}^3

$$K = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle$$

$$\bar{X} = \langle \langle x_1, \dots, x_g \rangle \rangle_K$$

$$\bar{Y} = \langle \langle y_1, \dots, y_g \rangle \rangle_K$$

$$\mathcal{G} = \{ h \in \mathcal{M} \mid h_{\#}(\bar{X}) \subset \bar{X}, \quad h_{\#}(\bar{Y}) \subset \bar{Y} \}$$

Some related groups:

Handlebody groups


$$\mathcal{H} = \text{MCG}(V, D) = \{ h \in \mathcal{M} \mid h_{\#}(\bar{X}) \subset \bar{X} \}$$

$$\mathcal{H}' = \text{MCG}(V', D) = \{ h \in \mathcal{M} \mid h_{\#}(\bar{Y}) \subset \bar{Y} \}$$

Twist (or Luft) groups


$$\mathcal{T} = \ker(\mathcal{H} \longrightarrow \text{Aut}(\pi_1(V, *)))$$

$$\mathcal{T}' = \ker(\mathcal{H}' \longrightarrow \text{Aut}(\pi_1(V', *)))$$




- **finitely generated**
- **not finitely generated**

Examples



- $t_\epsilon, t_\gamma, t_\delta \in \mathcal{T}$
- $t_\epsilon, t_\gamma \notin \mathcal{G}$
- $t_\epsilon t_\gamma^{-1}, t_\delta \in \mathcal{G}$




Action on $H_1(\Sigma, \mathbb{Z})$

$$K = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle$$

$$a_i = [x_i], \quad b_i = [y_i]$$

$$H = H_1(\Sigma; \mathbb{Z}) = \langle a_1, b_1, \dots, a_g, b_g \rangle_{\text{ab}}$$

$$\text{Sp}(H, \omega) \simeq \text{Sp}(2g, \mathbb{Z})$$



$$\omega : H \otimes H \longrightarrow \mathbb{Z}$$

intersection form

Lemma (folklore)

Let $\sigma : \mathcal{M} \rightarrow \text{Sp}(2g, \mathbb{Z})$ be the action on H , then

- $\sigma(\mathcal{H}) = \left\{ \begin{pmatrix} P & R \\ 0 & (P^T)^{-1} \end{pmatrix} \mid P^{-1}R \text{ is symmetric} \right\}$
- $\sigma(\mathcal{H}') = \left\{ \begin{pmatrix} P & 0 \\ R & (P^T)^{-1} \end{pmatrix} \mid P^T R \text{ is symmetric} \right\}$
- $\sigma(\mathcal{G}) = \left\{ \begin{pmatrix} P & 0 \\ 0 & (P^T)^{-1} \end{pmatrix} \mid P \in \text{GL}(g, \mathbb{Z}) \right\} \simeq \text{GL}(g, \mathbb{Z})$
- $\sigma(\mathcal{T}) = \left\{ \begin{pmatrix} \text{Id}_g & R \\ 0 & \text{Id}_g \end{pmatrix} \mid R \text{ is symmetric} \right\} \simeq \mathbb{Z}^{\frac{1}{2}g(g+1)}$
- $\sigma(\mathcal{T}') = \left\{ \begin{pmatrix} \text{Id}_g & 0 \\ R & \text{Id}_g \end{pmatrix} \mid R \text{ is symmetric} \right\} \simeq \mathbb{Z}^{\frac{1}{2}g(g+1)}$

Double Johnson filtration for the Goeritz group

$$K = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle \quad \bar{X} = \langle\langle x_1, \dots, x_g \rangle\rangle_K \quad \bar{Y} = \langle\langle y_1, \dots, y_g \rangle\rangle_K$$

The Goeritz group

$$\mathcal{G} = \{ h \in \mathcal{M} \mid h_{\#}(\bar{X}) \subset \bar{X}, \quad h_{\#}(\bar{Y}) \subset \bar{Y} \}$$

acts on K preserving the double lower central series $(K_{i,j})_{(i,j) \in \mathbb{N}^2}$ of $(K; \bar{X}, \bar{Y})$

For all $(m, n) \in \mathbb{N}^2$ we set

$$\mathcal{G}_{m,n} = \{ h \in \mathcal{G} \mid [h, K_{i,j}] \subset K_{m+i, n+j} \text{ for all } (i, j) \in \mathbb{N}^2 \} \quad [h, z] = h_{\#}(z)z^{-1}$$

Proposition

- $\mathcal{G}_{m,n} = \{ h \in \mathcal{G} \mid [h, \bar{X}] \subset K_{m+1, n}, [h, \bar{Y}] \subset K_{m, n+1} \}$
- $\mathcal{G}_{0,0} = \mathcal{G}$
- $\mathcal{G}_{m,n} \trianglelefteq \mathcal{G}$
- $[\mathcal{G}_{m,n}, \mathcal{G}_{a,b}] \subset \mathcal{G}_{m+a, n+b}$ that is, $(\mathcal{G}_{m,n})_{m,n}$ is a \mathbb{N}^2 -filtration.

Examples

$$t_\epsilon t_\gamma^{-1} \in \mathcal{G}_{1,0}$$

$$t_\delta \in \mathcal{G}_{1,1}$$


$$\begin{matrix} \mathcal{G}_{0,0} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{0,1} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{0,2} \\ \cup \end{matrix} \supset \cdots$$

$$\begin{matrix} \mathcal{G}_{1,0} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{1,1} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{1,2} \\ \cup \end{matrix} \supset \cdots$$

$$\begin{matrix} \mathcal{G}_{2,0} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{2,1} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{2,2} \\ \cup \end{matrix} \supset \cdots$$

$$\begin{matrix} \mathcal{G}_{3,0} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{3,1} \\ \cup \end{matrix} \supset \begin{matrix} \mathcal{G}_{3,2} \\ \cup \end{matrix} \supset \cdots$$

$$\vdots \qquad \vdots \qquad \vdots$$



Double Johnson filtration for the Mapping class group

$$K = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle \quad \bar{X} = \langle \langle x_1, \dots, x_g \rangle \rangle_K \quad \bar{Y} = \langle \langle y_1, \dots, y_g \rangle \rangle_K$$
$$(K_{i,j})_{(i,j) \in \mathbb{N}^2}: \text{ double lower central series of } (K; \bar{X}, \bar{Y})$$

We extend the definition of
 $(K_{i,j})_{(i,j) \in \mathbb{N}^2}$ as follows:

$\forall (i,j) \in \mathbb{Z}^2$, set

$$K_{i,j} := K_{\max(0,i), \max(0,j)}$$

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \vdots & \\ \cdots & K_{0,0} & \supset & K_{0,0} & \supset & K_{0,1} & \supset K_{0,2} \cdots \\ & \cup & & \cup & & \cup & \cup \\ \cdots & K_{0,0} & \supset & K_{0,0} & \supset & K_{0,1} & \supset K_{0,2} \cdots \\ & \cup & & \cup & & \cup & \cup \\ \cdots & K_{1,0} & \supset & K_{1,0} & \supset & K_{1,1} & \supset K_{1,2} \cdots \\ & \cup & & \cup & & \cup & \cup \\ \cdots & K_{2,0} & \supset & K_{2,0} & \supset & K_{2,1} & \supset K_{2,2} \cdots \\ & \cup & & \cup & & \cup & \cup \\ \cdots & K_{3,0} & \supset & K_{3,0} & \supset & K_{3,1} & \supset K_{3,2} \cdots \\ & \cup & & \cup & & \cup & \cup \\ & \vdots & & \vdots & & \vdots & \vdots \end{array}$$

Double Johnson filtration for the Mapping class group

For $(m, n) \in \mathbb{Z}^2$, we set


$$\mathcal{M}_{m,n} := \{ h \in \mathcal{M} \mid [h^{\pm 1}, K_{i,j}] \subset K_{m+i, m+j} \quad \forall (i,j) \in \mathbb{N}^2 \} \leq \mathcal{M}$$

Theorem

For $(m, n) \in \mathbb{Z}^2$, we have

- $\mathcal{M}_{m,n} = \{ h \in \mathcal{M} \mid [h^{\pm 1}, K_{1,0}] \subset K_{m+1,n} \text{ and } [h^{\pm 1}, K_{0,1}] \subset K_{m,n+1} \}$
- $\mathcal{M}_{m,n} = \mathcal{M}_{\max(-1,m), \max(-1,n)}$
- If $(m, n) \in \mathbb{N}^2$, then $\mathcal{M}_{m,n} = \mathcal{G}_{m,n}$
- $\mathcal{M}_{1,-1} = \mathcal{T}$ and $\mathcal{M}_{-1,1} = \mathcal{T}'$ (twist groups)
- $\mathcal{M}_{0,-1} = \mathcal{H}$ and $\mathcal{M}_{-1,0} = \mathcal{H}'$ (handlebody groups)
- $\mathcal{M}_{-1,-1} = \mathcal{M}$ (mapping class group)

Double Johnson filtration for the Mapping class group



Proposition

For $(m, n), (a, b) \in \{(k, l) \in \mathbb{Z}^2 \mid k, l \geq -1, k + l \geq 1\} \cup \{(0, 0)\}$, we have


$$[\mathcal{M}_{m,n}, \mathcal{M}_{a,b}] \subset \mathcal{M}_{m+a, n+b}$$

Examples


D_n° be a disk with n holes

$\text{FPB}_n = \text{MCG}(D_n^\circ)$ framed pure braid group on n strands


Any embedding of $D_n^\circ \rightarrow \Sigma$ induces a map $\text{FPB}_n \rightarrow \mathcal{M}$



$$D_{2g}^\circ \longrightarrow \Sigma$$



$$D_g^\circ \longrightarrow \Sigma$$



$$D_g^\circ \longrightarrow \Sigma$$

Proposition

Consider an embedding $D_n^\circ \rightarrow \Sigma$ such that the image of the holes of D_n° bound mutually disjoint disks in V . Let $f : \text{FPB}_n \rightarrow \mathcal{M}$ be the induced homomorphism. If $\beta \in \Gamma_k \text{FPB}_n$, then $f(\beta) \in \mathcal{M}_{k,-1}$.

Relation with the usual Johnson filtration

$$K = \pi_1(\Sigma, *)$$

$$\underbrace{\mathcal{I} = J_1 \mathcal{M}}_{\text{Torelli}} \supset \underbrace{\mathcal{K} = J_2 \mathcal{M}}_{\text{Johnson}} \supset J_3 \mathcal{M} \supset \dots$$

$$J_n \mathcal{M} = \{h \in \mathcal{M} \mid [h, K] \subset \Gamma_{n+1} K\}$$

$\forall m, n \geq -1$ with $m + n \geq 1$ we have

$$\mathcal{M}_{m,n} \subset J_{m+n} \mathcal{M}$$

\mathcal{M}	\mathcal{H}'	\mathcal{T}'	$\mathcal{M}_{-1,2}$	\dots
\mathcal{H}	\mathcal{G}	$\mathcal{M}_{0,1}$	$\mathcal{M}_{0,2}$	\dots
\mathcal{T}	$\mathcal{M}_{1,0}$	$\mathcal{M}_{1,1}$	$\mathcal{M}_{1,2}$	\dots
$\mathcal{M}_{2,-1}$	$\mathcal{M}_{2,0}$	$\mathcal{M}_{2,1}$	$\mathcal{M}_{2,2}$	\dots
$\mathcal{M}_{3,-1}$	$\mathcal{M}_{3,0}$	$\mathcal{M}_{3,1}$	$\mathcal{M}_{3,2}$	\dots
⋮	⋮	⋮	⋮	⋮

- $\mathcal{M}_{2,-1}, \mathcal{M}_{1,0}, \mathcal{M}_{0,1}, \mathcal{M}_{-1,2} \subset J_1 \mathcal{M} = \mathcal{I}$

- $\mathcal{M}_{3,-1}, \mathcal{M}_{2,0}, \mathcal{M}_{1,1}, \mathcal{M}_{0,2}, \mathcal{M}_{-1,3} \subset J_2 \mathcal{M} = \mathcal{K}$

Relation with the usual Johnson filtration

Theorem

$$\mathcal{I} = \mathcal{M}_{2,-1} \cdot \mathcal{M}_{1,0} \cdot \mathcal{M}_{0,1} \cdot \mathcal{M}_{-1,2} \cdot \mathcal{K}$$

Conjecture

We have $\mathcal{K} \subset \mathcal{M}_{2,-1} \cdot \mathcal{M}_{1,0} \cdot \mathcal{M}_{0,1} \cdot \mathcal{M}_{-1,2}$ and therefore

$$\mathcal{I} = \mathcal{M}_{2,-1} \cdot \mathcal{M}_{1,0} \cdot \mathcal{M}_{0,1} \cdot \mathcal{M}_{-1,2}.$$

Theorem (case of automorphisms of free groups)

Let $F = F_{p,q} = \langle x_1, \dots, x_p, y_1, \dots, y_q \rangle$ be the free group on $p+q$ generators.

We can define a double filtration $(\mathcal{A}_{i,j})_{i,j \geq -1}$ for $\text{Aut}(F)$.

If $\mathcal{IA} = \ker(\text{Aut}(F) \rightarrow \text{Aut}(F/[F,F]))$, then

$$\mathcal{IA} = \mathcal{A}_{2,-1} \cdot \mathcal{A}_{1,0} \cdot \mathcal{A}_{0,1} \cdot \mathcal{A}_{-1,2}.$$

Double Johnson homomorphisms

$$K = \langle x_1, \dots, x_g, y_1, \dots, y_g \rangle \quad \bar{X} = \langle \langle x_1, \dots, x_g \rangle \rangle_K \quad \bar{Y} = \langle \langle y_1, \dots, y_g \rangle \rangle_K$$

$(K_{i,j})_{(i,j) \in \mathbb{N}^2}$: double lower central series of $(K; \bar{X}, \bar{Y})$

$$a_i = [x_i], \quad b_i = [y_i]$$

$$\mathbb{N}_+^2 = \{(m, n) \in \mathbb{N}^2 \mid m + n \geq 1\}$$

$$H = \langle a_1, b_1, \dots, a_g, b_g \rangle_{ab} \quad A = \langle a_1, \dots, a_g \rangle_{ab} \quad B = \langle b_1, \dots, b_g \rangle_{ab}$$

$$\mathfrak{Lie}(A, B) = \bigoplus_{(m,n) \in \mathbb{N}_+^2} \mathfrak{Lie}_{m,n}(A, B) \quad \mathfrak{Lie}_{0,1}(A, B) \quad \mathfrak{Lie}_{0,2}(A, B) \cdots$$

$$\\ \mathfrak{Lie}_{1,0}(A, B) \quad \mathfrak{Lie}_{1,1}(A, B) \quad \mathfrak{Lie}_{1,2}(A, B) \cdots$$

$$\begin{array}{llll} \mathbb{N}_+^2\text{-graded free Lie algebra} & \mathfrak{Lie}_{2,0}(A, B) & \mathfrak{Lie}_{2,1}(A, B) & \mathfrak{Lie}_{2,2}(A, B) \cdots \\ \text{generated by } A \text{ in degree } (1, 0) & \mathfrak{Lie}_{3,0}(A, B) & \mathfrak{Lie}_{3,1}(A, B) & \mathfrak{Lie}_{3,2}(A, B) \cdots \\ \text{and } B \text{ in degree } (0, 1). & \vdots & \vdots & \vdots \end{array}$$

Lemma

$$\bigoplus_{(m,n) \in \mathbb{N}_+^2} \frac{K_{m,n}}{K_{m+1,n} \cdot K_{m,n+1}} \simeq \bigoplus_{(m,n) \in \mathbb{N}_+^2} \mathfrak{Lie}_{m,n}(A, B) = \mathfrak{Lie}(A, B)$$

Proposition

For $(m, n) \in \mathbb{N}_+^2$ there is a subgroup

$$D_{m,n}(A, B) \leq (A \otimes \mathfrak{Lie}_{m,n+1}(A, B)) \oplus (B \otimes \mathfrak{Lie}_{m+1,n}(A, B))$$

and group homomorphisms

$$\tau_{m,n} : \mathcal{M}_{m,n} \longrightarrow D_{m,n}(A, B)$$

such that

$$\mathcal{M}_{m+1,n} \cdot \mathcal{M}_{m,n+1} \subset \ker(\tau_{m,n}).$$

Moreover, these homomorphisms are compatible with the usual Johnson homomorphisms: the diagram

$$\begin{array}{ccc} \mathcal{M}_{m,n} & \xrightarrow{\subseteq} & J_{m+n}\mathcal{M} \\ \tau_{m,n} \downarrow & & \downarrow \tau_{m+n} \\ D_{m,n}(A, B) & \xrightarrow{j} & D_{m+n}(H) \end{array}$$

is commutative.

Remark

The above result can be extended to all the double filtration

$$\left(\tau_{m,n} : \mathcal{M}_{m,n} \longrightarrow D_{m,n}(A, B) \right)_{m,n \geq -1}$$

We also obtain the following: If $h \in \mathcal{I}$, then there exist

$$h_1 \in \mathcal{M}_{2,-1}, \quad h_2 \in \mathcal{M}_{1,0}, \quad h_3 \in \mathcal{M}_{0,1}, \quad h_4 \in \mathcal{M}_{-1,2}$$

such that

$$\tau_1(h) = \tau_{2,-1}(h_1) + \tau_{1,0}(h_2) + \tau_{0,1}(h_3) + \tau_{-1,2}(h_4).$$